

AD-A191 892

CONVERGENCE RATE OF CODES FOR NUMERICAL QUADRATURE  
TECHNIQUES FOR CLASSICAL RAY TRACING(U) NAVAL OCEAN  
SYSTEMS CENTER SAN DIEGO CA E R FLOYD DEC 86


1/1

UNCLASSIFIED

F/G 20/1

NL





DTIC FILE COPY

AD-A191 892

DTIC  
REPORT DOCUMENTATION PAGE  
ELECTE

MAR 1 7 1988

2a SECURITY CLASSIFICATION AUTHORITY		1b RESTRICTIVE MARKINGS	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
4 PERFORMING ORGANIZATION REPORT NUMBER(S)		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION Naval Ocean Systems Center	6b OFFICE SYMBOL (if applicable)	7a NAME OF MONITORING ORGANIZATION	
6c ADDRESS (City, State and ZIP Code) San Diego, CA 92152-5000		7b ADDRESS (City, State and ZIP Code)	
8a NAME OF FUNDING/SPONSORING ORGANIZATION Commander Sub Force Pacific	8b OFFICE SYMBOL (if applicable) CSFP	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c ADDRESS (City, State and ZIP Code) U.S. Pacific Fleet Pearl Harbor, HI 96860		10 SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO DMN	
		PROJECT NO MR02	TASK NO SUBPAC
		AGENCY ACCESSION NO DN088 580	
11 TITLE (Include Security Classification) Convergence Rate of Codes for Numerical Quadrature Techniques for Classical Ray Tracing			
12 PERSONAL AUTHOR(S) E.R. Floyd			
13a TYPE OF REPORT Presentation	13b TIME COVERED FROM TO	14 DATE OF REPORT (Year, Month, Day) December 1986	15 PAGE COUNT
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES FIELD GROUP SUB GROUP		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) raytracing Stieltjes measure	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) The estimated residual error (or its bound) for numerical quadratures is usually expressed in terms of a derivative of some order of the integrand or some residual factor of the integrand after factoring out a countable number of zeroes and singularities that occur along the integration path. The order of the derivative is a function of the number of sample points for evaluating the integrand. Regrettably, the magnitudes of these higher order derivatives are difficult enough to estimate for even analytic sound velocity profiles. In practice, observed sound velocity profiles, which are usually given in tabular form and include measurement errors, exacerbate our inability to assess the magnitudes of these higher order derivatives. An estimate of the residual error expressed in terms of a first derivative would be far more practical for both analytic and observed sound velocity profiles.			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL E.R. Floyd		22b TELEPHONE (Include Area Code) 619-225-6851	22c OFFICE SYMBOL Code 192

DD FORM 1473, 84 JAN

83 APR EDITION MAY BE USED UNTIL EXHAUSTED  
ALL OTHER EDITIONS ARE OBSOLETE


UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE

88 3 16 084


I8.

CONVERGENCE RATE OF CODES  
FOR NUMERICAL QUADRATURE TECHNIQUES  
FOR CLASSICAL RAY TRACING

Edward R. Floyd  
Arctic Submarine Laboratory



By classical ray tracing, determine range over a single Riemann sheet for a ray described by the constant of motion (vertex velocity)  $C_m$ . How accurate?



Accession For

NTIS GRA&I ☒  
 DTIC TAB ☐  
 Unannounced ☐  
 Justification

By  
 Distribution/

Availability Codes

Dist	Avail and/or Special
A-1	

Range Integral:

$$\begin{aligned} R_s &= \int_{z_0}^{z_v} \frac{C(z)}{[C_m^2 - C^2(z)]^{1/2}} dz \\ &= \int_{z_0}^{z_v} W(z) f(z) dz \end{aligned}$$

$W$  and  $f$  are convenient functions of  $z$ .  $[W dz]$  becomes a Stieltjes measure and  $f(z)$ , the Stieltjes integrand.

Trick: Choose  $W(z)$  to incorporate any poor behavior (singularities or zeros) into  $W$  leaving  $f(z)$  well behaved.

Thus the Gaussian quadrature

$$R_s = (z_V - z_0)^m \sum_{i=1}^n w_{i,n} f(z_{i,n})$$

where:

$n$  is the order of the approximation;

$m$  is determined by  $W$ ;

$w_{i,n}$  is a Gaussian weighting coefficient  
determined by  $W$ ;

$z_{i,n}$  is a Gaussian sample point determined  
by  $W$ .

The Gaussian error is given by

$$\eta_n = \frac{f^{(2n)}(\xi)}{(2n)!} \int_{z_0}^{z_v} W(z) \left[ \prod_{i=1}^n (z - z_{i,n}) \right]^2 dz$$

$$\text{where } z_0 \leq \xi \leq z_v.$$

For underwater acoustics

$$W = (z_v - z)^{-1/2}.$$

Factors out branch point singularity at vertex depth.

Therefore

$$\eta_n = \frac{\pi^{1/2} (z_v - z_0)^{2n+1}}{4n+1} \frac{[(2n)!]^2}{[(4n)!]^2} f^{(2n)}(\xi),$$

$$z_0 \leq \xi \leq z_v.$$



$$\eta_n = \frac{\pi^{1/2} (z_v - z_0)^{2n+1}}{4n+1} \frac{[(2n)!]^3}{[(4n)!]^2} f^{(2n)}(\xi),$$

$$z_0 \leq \xi \leq z_v.$$

---

This expression has several deficiencies;

- Usually impractical to evaluate  $f^{(2n)}$ .
- $f^{(i)}$  for  $i < 2n$  may be discontinuous.
- $C(z)$  is often known only in tabular form.

$$\eta_n = \frac{\pi^{1/2} (z_v - z_0)^{2n+1}}{4n+1} \frac{[(2n)!]^3}{[(4n)!]^2} f^{(2n)}(\xi),$$

$$z_0 \leq \xi \leq z_v.$$

---

Analytic (hypothetical)  $C(z)$ :\*

- Rapid convergence with increasing  $n$ .
- Only limited by round-off error.
- Thumb rule for maximum effective  $n$ :

$$n_{\max} \approx (\text{"number of significant figures"})/2$$


---

\* E. R. Floyd, J. Acoust. Soc. Am. 49, 1580-1590 (1971)

$$\eta_n = \frac{\pi^{1/2} (z_V - z_0)^{2n+1}}{4n+1} \frac{[(2n)!]^3}{[(4n)!]^2} f^{(2n)}(\xi),$$

$$z_0 \leq \xi \leq z_V.$$

---

Observed (nonanalytic)  $C(z)$ :\*

- $C(z)$  tabled at discrete depths.
- Cubic spline fit  $\longrightarrow d^3C/dz^3$  discontinuous.  
Nevertheless still rapidly convergent over an assembly of 116 observed  $C(z)$ 's. MOE ---  $\eta$  reduced by a factor of 0.763 for each additional sample point per Riemann sheet.
- Thumb rule for maximum effective  $n$ :

$$\eta_{\max} \geq \text{"number of significant figures"}.$$

---

\* E. R. Floyd and J. D. Pugh, J. Acoust. Soc. Am. 61, 682-687 (1977).

Present Status:

- Even for analytic functions, quadrature errors are difficult to estimate because higher order derivatives are generally complicated.
- While we have convergence rates for an assembly of observed profiles, the discontinuity of higher order derivatives confounds our estimate of convergence for a particular observed profile.
- Desire an estimate expressed in terms of the first derivative of the Stieltjes integrand.

END

DATE

FILMED

6-88

DTIC